
system
e
cluster
mass
typical
stellar mass
event
Supernova
wind
km
excess
log
ed
g
progenitor
velocity
target
rest frame
B
Astrophysical
model
M
thus
number
C rich
higher
de
best fit
properties
rate
timescale
evolution
redshift
transient
period
cluster members
early
irradiation
BlackHole
count rate
wavelength
Xray
peak
time
region
radius
emission line
photometric
prior
hydrogen
uncertainties
f
component
background
difference
analysis
NICER
around
Among
best
phase
t
host
temperature
outburst
Supernovae
color
scale
ultra stripped
optical
well
UV
absorption
small
Mira
energy
measurement
emission
sample
O rich
strong
red
normal
follow uncertainty
point
result
distribution
II
radio
step
power law
Swift
blackbody
low
survey
distance
SEDM
MJD
catalog
u
Telescope
photometry
ratio
ZTF
low mass
bright
explosion
Sky Survey
density
blue
case
light curve
host galaxy
keV
detection
Galactic
NGC
consistent
Note
discovery
observed
band
template
relation
mass star
shock
Supernovae Ia
SED
function
position
method
total
NuSTAR
erg
stellar
type
median
epoch
high
optical
disk
uv
candidate
power
photon
less
magnitude
luminosity
disk frequency
TidalDisruptionEvent
early time
feature different
optical emission